



Local exact controllability to the trajectories of the Korteweg–de Vries–Burgers equation on a bounded domain with mixed boundary conditions

Eduardo Cerpa^a, Cristhian Montoya^{a,*}, Bingyu Zhang^b

^a Departamento de Matemática, Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile

^b Department of Mathematical Sciences, University of Cincinnati, Cincinnati, OH 45221, USA

Received 28 February 2019; revised 31 July 2019; accepted 29 October 2019

Abstract

This paper studies the internal control of the Korteweg–de Vries–Burgers (KdVB) equation on a bounded domain. The diffusion coefficient is time-dependent and the boundary conditions are mixed in the sense that homogeneous Dirichlet and periodic Neumann boundary conditions are considered. The exact controllability to the trajectories is proven for a linearized system by using duality and getting a new Carleman estimate. Then, using an inversion theorem we deduce the local exact controllability to the trajectories for the original KdVB equation, which is nonlinear.

© 2019 Elsevier Inc. All rights reserved.

Keywords: Korteweg–de Vries–Burgers equation; Controllability; Carleman estimates

1. Introduction

The Korteweg–de Vries (KdV) equation appears in the nineteenth century with the works of Boussinesq [5], Korteweg and de Vries [24], [29]. From a physical point of view, the KdV equation represents a model for the motion of long water waves in channels of shallow depth, in which two different phenomena are presented, namely, nonlinear convection and dispersion. This

* Corresponding author.

E-mail addresses: eduardo.cerpa@usm.cl (E. Cerpa), cristhian.montoya@usm.cl (C. Montoya), zhangb@ucmail.uc.edu (B. Zhang).

<https://doi.org/10.1016/j.jde.2019.10.043>

0022-0396/© 2019 Elsevier Inc. All rights reserved.

interaction produces a wave traveling at constant speed without losing its sharp, usually called *soliton*.

The study of the KdV equation from a control point of view began with the work of Russell [33] and Zhang [35] in late 1980s. Both exact control problem and stabilization problem have been intensively studied since then. For internal control of the KdV equation on a periodic domain, Russell and Zhang [34] showed that the system is locally exactly controllable and exponentially stabilizable in the space $H^s(\mathbb{T})$ for any $s \geq 0$. Their work was improved by Laurent, Rosier and Zhang [25] who showed that the system is globally exponentially stabilizable and (large time) globally exactly controllable in $H^s(\mathbb{T})$ for any $s \geq 0$. The study of the boundary controllability for the KdV equation on a bounded domain $(0, L)$ was started by Rosier [31] where he employed only one control input. Using compactness–uniqueness arguments and the Hilbert Uniqueness method he first showed surprisingly that the linearized system around the origin is exact controllable in the space $L^2(0, L)$ if and only if the length L of the spatial domain does not belong to a set of critical values. Then assuming the length L of the spatial domain is not critical, he showed the nonlinear system is locally exactly controllable in the space $L^2(0, L)$ by using contraction mapping principle. If all three boundary controls are employed, Zhang [36] using a different approach proved that the system is locally exactly controllable in $H^s(0, L)$ for $s \geq 0$ without any restrictions on the spatial domain. When the linearized system is not controllable, nevertheless, one can still prove that the nonlinear system is locally exact controllable in the space $L^2(0, L)$ by using power series expansion of the solutions (see [13,9,10]). Other related results can be found in [19] and [32]. Concerning the internal controllability for the KdV equation on a bounded domain with homogeneous Dirichlet boundary conditions, the most recent work was done by Capistrano–Filho et al. in [7], where the authors obtained some controllability results using an approach based on Carleman estimates and weighted Sobolev spaces.

On the other side, the Burgers equation first appeared in 1940 as a simplified one-dimensional model for the Navier–Stokes system [6]. Its controllability properties on bounded domains are certainly different in each case (i.e., distributed controls, boundary control, initial value control). For instance, in [22], Horsin studies the exact controllability on a bounded domain for the Burgers equation by means of the return method [12]. In the case of boundary controllability with partial measurements, the work [20] done by Imanuvilov and Guerrero shows that the exact controllability property does not hold. In the context of distributed controls with Dirichlet and Neumann boundary conditions, the works by Fernández–Cara and Guerrero [15] and Marbach [28] addressed these problems.

As consequence of the union of the KdV and Burgers equations arise the Korteweg–de Vries–Burgers equation (KdVB equation), which in our case has homogeneous Dirichlet boundary conditions and periodic Neumann boundary conditions. More precisely, we consider the following system

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} + yy_x = F(x, t) & \text{in } (0, L) \times (0, T), \\ y(0, t) = y(L, t) = 0 & \text{in } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L), \end{cases} \quad (1)$$

where $y = y(x, t)$ represents the surface elevation of the water wave at time $(0, T)$ and space $(0, L)$, $\mu(t) := \mu_0 + \tilde{\mu}(t) > 0$, with $\mu_0 > 0$ and $\tilde{\mu}(t) \geq 0$ is the diffusion coefficient, $F = F(x, t)$ is an internal force and y_0 is the initial datum. The system (1) can be viewed as a model of propagation of long water waves in channels of shallow depth, whose solutions depend on the

nonlinearity, dispersion, and dissipation. Moreover, by introducing a variable coefficient $\mu(t)$, the KdVB equation (1) is useful to describe cosmic plasmas phenomena [18], [27]. Respect to the boundary conditions, they appear in order to make symmetric the operator. Thus, studying the controllability of our system can help to build for instance some feedback laws requiring that the underlying operator is skew-adjoint. Besides, we can explicitly mention the difficulty appearing with these boundary conditions: the hidden regularity $L^2(0, T; H^1(0, L))$ is not implied by the third order term. That is the reason that the Laplacian is added.

From a mathematical point of view, there exist several results for the KdVB equation in both bounded and unbounded domains, concerning the global and local well-posedness problem [8], [26], [14] and [3]; the optimal control problem [4], [11]; the internal controllability problem on unbounded domain [17]; and the boundary feedback stabilization problem [23]. As far as we know, the internal controllability problem for (1) has not been studied and thus, our paper will fill this gap.

Throughout our work, we will use the following notation: let $\omega \subset (0, L)$ be a nonempty open subset and let $Q = (0, L) \times (0, T)$, for $T > 0$. The main result of this paper is related to the local exact controllability to the trajectories of the KdVB equation

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} + yy_x = v1_{\omega \times (0, T)} & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L), \end{cases} \quad (2)$$

where $v = v(x, t)$ stands for the control, which acts in the domain $\omega \times (0, T)$.

Let us now introduce the concept of *exact controllability to the trajectories* for the Korteweg–de Vries–Burgers (KdVB) equation. The goal is to reach (in finite time T) any point on a given trajectory of the same operator. Let $\bar{y}$ be a solution of the uncontrolled KdVB equation:

$$\begin{cases} \bar{y}_t + \bar{y}_{xxx} - \mu(t)\bar{y}_{xx} + \bar{y}\bar{y}_x = 0 & \text{in } Q, \\ \bar{y}(0, t) = \bar{y}(L, t) = 0 & \text{on } (0, T), \\ \bar{y}_x(0, t) = \bar{y}_x(L, t) & \text{on } (0, T), \\ \bar{y}(\cdot, 0) = \bar{y}_0(\cdot) & \text{in } (0, L). \end{cases} \quad (3)$$

We look for a control v such that the solution of (2) satisfies:

$$y(\cdot, T) = \bar{y}(\cdot, T) \quad \text{in } (0, L). \quad (4)$$

In this paper we will show that for any given trajectory $\bar{y}$, which is a solution of (3), there exists a $\delta > 0$ such that, for any $y_0 \in X$ (an appropriate Banach space) satisfying

$$\|y_0 - \bar{y}_0\|_X \leq \delta, \quad (5)$$

one can find a control v such that the system (2) admits a solution $y(x, t)$ satisfying (4).

Here we assume

$$\bar{y} \in C([0, T]; H^s(0, L)) \cap L^2(0, T; H^{s+1}(0, L)) \quad (6)$$

for some $s \in [0, 3]$.

To prove the exact controllability to the trajectory, we consider two relevant control systems, namely, the linearized system of (2) around $\bar{y}$ which is

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} + \bar{y}y_x + y\bar{y}_x = f + v1_{\omega \times (0,T)} & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L) \end{cases} \quad (7)$$

and the adjoint system associated to (7)

$$\begin{cases} -\varphi_t - \varphi_{xxx} - \mu(t)\varphi_{xx} - \bar{y}\varphi_x = g & \text{in } Q, \\ \varphi(0, t) = \varphi(L, t) = 0 & \text{on } (0, T), \\ \varphi_x(0, t) = \varphi_x(L, t) & \text{on } (0, T), \\ \varphi(\cdot, T) = \varphi_T(\cdot) & \text{in } (0, L). \end{cases} \quad (8)$$

Our strategy is as follows:

- i) Establish first a global Carleman inequality for the system (8). More precisely, we will prove the following Theorem:

Theorem 1.1. *Let $\mu \in L^\infty(0, T)$ and assume that $\bar{y}$ satisfies (6). Then, there exist two positive constants s_0, C depending on L and ω such that, for every $\varphi_T \in L^2(0, L)$ and $g \in L^2(Q)$, the corresponding solution to (8) satisfies:*

$$\begin{aligned} & \iint_Q (s^5 \xi^5 |\varphi|^2 + s^3 \xi^3 |\varphi_x|^2 + s \xi |\varphi_{xx}|^2) e^{-4s\hat{\alpha}} dx dt \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\alpha}} dx dt + s^7 \iint_{\omega \times (0, T)} \xi^7 e^{-6s\hat{\alpha} + 2s\hat{\alpha}} |\varphi|^2 dx dt \right), \end{aligned} \quad (9)$$

for every $s \geq s_0$.

The estimate (9) allows us to prove a null controllability result for the linear system (7) with right-hand side satisfying suitable decreasing properties near $t = T$. Theorem 1.1 will be proved using the same approach as in [21, 2, 7].

- ii) Then establish the local exact controllability to the trajectories for the KdVB equation. Here, fixed point arguments will be used to prove Theorem 1.2 given below.

Theorem 1.2. *Let $T > 0$ be given. Assume $\bar{y} \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))$ is the solution of (3). Then there exists a $\delta > 0$ such that for $y_0 \in L^2(0, L)$ satisfying (5), one can find a function control $v \in L^2(0, T; L^2(\omega))$ such that (2) admits a solution y satisfies*

$$y(\cdot, T) = \bar{y}(\cdot, T) \quad \text{in } (0, L).$$

The paper is organized as follows. In Section 2, we prove the local well-posedness of the system (1). In Section 3, we establish a Carleman inequality for the adjoint system (8), which is

associated to the linearized KdVB equation. In other words, we prove Theorem 1.1. In Section 4, we deal with the null controllability for a linearized system with a right-hand side in $L^2(0, L)$. Finally, in Section 5, the proof of Theorem 1.2 is given.

2. Well-posedness

2.1. Linear case

In this subsection we establish the well-posedness of the system

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} + \bar{y}y_x + \bar{y}_x y = f & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L), \end{cases} \quad (10)$$

where $\bar{y}$ satisfies (3). First we consider the following linear problem

$$\begin{cases} y_t + y_{xxx} - \mu_0 y_{xx} = f & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L), \end{cases} \quad (11)$$

where $\mu_0 > 0$ is a constant.

Proposition 2.1. *Let $T > 0$ be given. For any $y_0 \in L^2(0, L)$ and $f \in L^1(0, T; L^2(0, L))$, (11) admits a unique mild solution $y \in C([0, T]; L^2(0, L))$ satisfying*

$$\|y\|_{C([0, T]; L^2(0, L))} \leq C(\|y_0\|_{L^2(0, L)} + \|f\|_{L^1(0, T; L^2(0, L))})$$

where $C > 0$ is a constant independent of y_0 and f .

Proof. Consider the operator $A := -\partial_x^3 + \mu_0 \partial_x^2$ defined on

$$\mathcal{D}(A) := \{u \in H^3(0, L) \cap H_0^1(0, L) : u(0) = u(L) = 0, u_x(0) = u_x(L)\} \subset L^2(0, L).$$

For any $\varphi \in \mathcal{D}(A)$,

$$\langle A\varphi, \varphi \rangle_{L^2(0, L)} = - \int_0^L \varphi_{xxx} \varphi \, dx + \mu_0 \int_0^L \varphi_{xx} \varphi \, dx = -\mu_0 \int_0^L |\varphi_x|^2 \, dx \leq 0.$$

Thus A is dissipative. Similarly, one can verify that A^* is also dissipative. Thus, the operator A generates a strongly semigroup $\{S(t)\}_{t \geq 0}$ of contractions in $L^2(0, L)$ by the Lumer–Phillips Theorem (see [30], Corollary 4.4, page 15). Hence, for any $y_0 \in L^2(0, L)$, $T > 0$ and $f \in L^1(0, T; L^2(0, L))$, (11) admits a unique mild solution $y \in C([0, T]; L^2(0, L))$, given by the formula

$$y(t) = S(t)y_0 + \int_0^t S(t-s)f(s)ds, \quad \forall t \in [0, T] \quad (12)$$

and depending continuously on the data, i.e.,

$$\|y\|_{C([0,T];L^2(0,L))} := \sup_{t \in [0,T]} \|y\|_{L^2(0,L)} \leq (\|y_0\|_{L^2(0,L)} + \|f\|_{L^1(0,T;L^2(0,L))}).$$

This completes the proof of Proposition 2.1. $\square$

Remark 2.1. Observe that if the initial data y_0 belongs to $\mathcal{D}(A)$ and $f \in C^1([0, T]; L^2(0, L))$ or $f \in L^1(0, T; \mathcal{D}(A)) \cap C([0, T]; L^2(0, L))$, the system (11) admits a unique classical solution, in other words, y belongs to

$$C([0, T]; L^2(0, L)) \cap C^1((0, T]; L^2(0, L)) \cap C((0, T]; \mathcal{D}(A)),$$

which can be expressed as (12). The reader interested can see [30, Corollary 2.2, page 106] for more details.

The following lemma reveals a global Kato smoothing property of the mild solutions of (11).

Lemma 2.1. *For every $T > 0$, $f \in L^1(0, T; L^2(0, L))$ and $y_0 \in L^2(0, L)$, the corresponding mild solution of (11) belongs to $C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))$ and satisfies*

$$\|y\|_{L^\infty(0,T;L^2(0,L))} + \|y\|_{L^2(0,T;H^1(0,L))} \leq C(\|y_0\|_{L^2(0,L)} + \|f\|_{L^1(0,T;L^2(0,L))}),$$

for some positive constant C dependent of μ_0 . Furthermore, the term yy_x belongs to $L^1(0, T; L^2(0, L))$ and it satisfies the estimate

$$\|yy_x\|_{L^1(0,T;L^2(0,L))} \leq C\|y\|_{L^2(0,T;H^1(0,L))}^2,$$

for some constant $C > 0$ dependent of μ_0 .

Proof. The proof follows the same ideas of [23], it is therefore omitted here. $\square$

Now we recall three additional Lemmas on sharp Kato smoothing property of the linear KdVB systems, which can be found in [23] as Lemma 2.7, Lemma 2.8 and Lemma 2.9.

The first one is for the linear KdVB equation posed on the whole line $\mathbb{R}$.

$$\begin{cases} w_t + w_{xxx} - \mu_0 w_{xx} = 0, & x \in \mathbb{R}, t \in (0, +\infty), \\ w(x, 0) = w_0(x), & x \in \mathbb{R}. \end{cases} \quad (13)$$

Lemma 2.2. *For a given $0 \leq s \leq 3$ and $w_0 \in H^s(\mathbb{R})$, the solution of problem (13) satisfies*

$$\sup_{x \in \mathbb{R}} \left(\|w(x, \cdot)\|_{H^{\frac{s+1}{3}}(0,+\infty)} + \|w_x(x, \cdot)\|_{H^{\frac{s}{3}}(0,+\infty)} \right) \leq C\|w_0\|_{H^s(\mathbb{R})}, \quad (14)$$

for some positive constant C .

The second one is for solutions of system (11).

Lemma 2.3. For given $y_0 \in L^2(0, L)$ and $f \equiv 0$, the unique solution y of (11) belongs to $L^\infty(0, L; H^{\frac{1}{3}}(0, T))$ with $y_x \in L^\infty(0, T; L^2(0, L))$ satisfying

$$\sup_{x \in [0, L]} \left(\|y(x, \cdot)\|_{H^{\frac{1}{3}}(0, T)} + \|y_x(x, \cdot)\|_{L^2(0, T)} \right) \leq C \|y_0\|_{L^2(0, L)}, \quad (15)$$

where C is a positive constant.

The third one is for solutions of the following linear problem

$$\begin{cases} y_t + y_{xxx} - \mu_0 y_{xx} = f & \text{in } (0, L) \times (0, +\infty), \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = 0 & \text{in } (0, L). \end{cases} \quad (16)$$

Lemma 2.4. For any $T > 0$ and $f \in L^1(0, T; L^2(0, L))$, there exists a positive constant C such that the solution $y(x, t)$ of (16) satisfies

$$\sup_{x \in [0, L]} \left(\|y(x, \cdot)\|_{H^{\frac{1}{3}}(0, T)} + \|y_x(x, \cdot)\|_{L^2(0, T)} \right) \leq C \int_0^T \|f(\cdot, s)\|_{L^2(0, L)} ds.$$

Combining the previous results, we obtain the following Lemma for the linear system (11).

Lemma 2.5. For any $T > 0$, $f \in L^1_{loc}(0, +\infty; L^2(0, L))$ and $y_0 \in L^2(0, L)$, the linear problem (11) admits a unique solution

$$y \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L)) \cap L^\infty(0, L; H^{\frac{1}{3}}(0, T))$$

satisfying $y_x \in L^\infty(0, L; L^2(0, T))$. Furthermore, there exists a constant C independent of T , y_0 and f such that

$$\begin{aligned} \sup_{t \in [0, T]} \|y(\cdot, t)\|_{L^2(0, L)} + \|y\|_{L^2(0, T; H^1(0, L))} + \sup_{x \in [0, L]} \left(\|y(x, \cdot)\|_{H^{\frac{1}{3}}(0, T)} + \|y_x(x, \cdot)\|_{L^2(0, T)} \right) \\ \leq C \left(\|f\|_{L^1(0, T; L^2(0, L))} + \|y_0\|_{L^2(0, L)} \right). \end{aligned}$$

In order to build the necessary regularity which will be used later on, we introduce a weak formulation of (11) for $f \in L^2(0, T; H^{-1}(0, L))$.

Definition 2.1. For $(f, y_0) \in L^2(0, T; H^{-1}(0, L)) \times L^2(0, L)$ a function $y \in C([0, T]; L^2(0, L))$ is called a weak solution of (11) if it satisfies the following identity

$$\int_0^T \int_0^L y g dx dt + (y(T), \varphi_T)_{L^2(0, L)} = \int_0^T \langle f, \varphi \rangle_{H^{-1}(0, L) \times H_0^1(0, L)} dt + (y_0, \varphi(0))_{L^2(0, L)}, \quad (17)$$

for all $(g, \varphi_T) \in L^1(0, T; L^2(0, L)) \times L^2(0, L)$, where $\varphi = \varphi(g, \varphi_T)$ is the mild solution of

$$\begin{cases} -\varphi_t - \varphi_{xxx} - \mu_0 \varphi_{xx} = g & \text{in } (0, L) \times (0, +\infty), \\ \varphi(0, t) = \varphi(L, t) = 0 & \text{on } (0, T), \\ \varphi_x(0, t) = \varphi_x(L, t) & \text{on } (0, T), \\ \varphi(\cdot, T) = \varphi_T & \text{in } (0, L). \end{cases} \quad (18)$$

In the following proposition we prove a regularity result for (11) by considering the pair (f, y_0) belongs to $L^2(0, T; H^{s-1}(0, L)) \times H^s(0, L)$ for any given $s \in [0, 3]$.

Proposition 2.2. *Let $0 \leq s \leq 3$ be given. For any $(f, y_0) \in L^2(0, T; H^{s-1}(0, L)) \times H^s(0, L)$, the system (11) admits a unique weak solution $y \in C([0, T]; H^s(0, L)) \cap L^2(0, T; H^{s+1}(0, L))$ and, furthermore, there exists a positive constant C such that*

$$\|y\|_{L^2(0, T; H^{s+1}(0, L))} \leq C \left(\|f\|_{L^2(0, T; H^{s-1}(0, L))} + \|y_0\|_{H^s(0, L)} \right). \quad (19)$$

Proof. Consider the system

$$\frac{dy}{dt} = Ay + f, \quad y(0) = y_0$$

as defined in (11). Since A is the infinitesimal generator of a semigroup $S(t)$ in the space $L^2(0, L)$, it follows from the standard semigroup theory that

$$y_0 \in L^2(0, L), \quad f \in L^1(0, T; L^2(0, L)) \implies y \in C([0, T]; L^2(0, L))$$

and moreover, there exists a constant $C > 0$ such that

$$\|y\|_{C([0, T]; L^2(0, L))} \leq C \left(\|y_0\|_{L^2(0, L)} + \|f\|_{L^1(0, T; L^2(0, L))} \right).$$

In addition,

$$y_0 \in \mathcal{D}(A), \quad f \in L^1(0, T; \mathcal{D}(A)) \implies y \in C([0, T]; H^3(0, L))$$

and furthermore, there exists a constant $C > 0$ such that

$$\|y\|_{C([0, T]; H^3(0, L))} \leq C \left(\|y_0\|_{H^3(0, L)} + \|f\|_{L^1(0, T; H^3(0, L))} \right).$$

Taking into account that

$$\frac{d}{dt} \int_0^L y^2(x, t) dx + 2\mu_0 \int_0^L y_x^2(x, t) dx = 2 \int_0^L f(x, t) y(x, t) dx$$

for any $t \geq 0$, we arrive at

$$\int_0^L y^2(x, t) dx - \int_0^L y^2(x, 0) dx + 2\mu_0 \int_0^t \int_0^L y_x^2(x, t) dx dt = 2 \int_0^t \int_0^L f(x, t) y(x, t) dx dt,$$

which implies that

$$\|y\|_{L^2(0, T; H^1(0, L))} \leq C (\|y_0\|_{L^2(0, L)} + \|f\|_{L^2(0, T; H^{-1}(0, L))}).$$

Similarly, if we let $w = Ay$, then we have

$$\|w\|_{L^2(0, T; H^1(0, L))} \leq C (\|Ay_0\|_{L^2(0, L)} + \|Af\|_{L^2(0, T; H^{-1}(0, L))}),$$

which yields that

$$\|y\|_{L^2(0, T; H^4(0, L))} \leq C (\|y_0\|_{H^3(0, L)} + \|f\|_{L^2(0, T; H^2(0, L))}).$$

By interpolation arguments,

$$\|y\|_{L^2(0, T; H^{1+3\theta}(0, L))} \leq C (\|y_0\|_{H^{3\theta}(0, L)} + \|f\|_{L^2(0, T; H^{-1+3\theta}(0, L))}),$$

for $0 \leq \theta \leq 1$, or in equivalent form

$$\|y\|_{L^2(0, T; H^{1+s}(0, L))} \leq C (\|y_0\|_{H^s(0, L)} + \|f\|_{L^2(0, T; H^{s-1}(0, L))}),$$

for $0 \leq s \leq 3$. This completes the proof of Proposition 2.2. $\square$

Now, we extend the previous Proposition to the linearized system (10). For this purpose, let us introduce the space Y_T^s as follows: for any $0 \leq s \leq 3$ and any $T > 0$,

$$Y_T^s := C([0, T]; H^s(0, L)) \cap L^2(0, T; H^{s+1}(0, L)).$$

Lemma 2.6. *For given $0 \leq s \leq 3$ and $T > 0$, there exists a positive constant C such that*

$$\|(uw)_x\|_{L^2(0, T; H^{s-1}(0, L))} \leq C \|u\|_{Y_T^s} \|w\|_{Y_T^s} \quad (20)$$

and

$$\|\mu w_{xx}\|_{L^2(0, T; H^{s-1}(0, L))} \leq C \|\mu\|_{L^\infty(0, T)} \|w\|_{Y_T^s} \quad (21)$$

holds for any $u, w \in Y_T^s$ and $\mu \in L^\infty(0, T)$.

Proof.

i) The case $s = 0$. In this case, we have

$$\|uw\|_{L^2(Q)}^2 \leq \int_0^T \|u(\cdot, t)\|_{L^\infty(0, L)}^2 \int_0^L w^2(x, t) dx dt \leq \|u\|_{L^2(0, T; L^\infty(0, L))}^2 \|w\|_{C([0, T]; L^2(0, L))}^2.$$

Taking into account that $H^1(0, L) \hookrightarrow L^\infty(0, L)$, the inequality (20) is proved.

On the other hand,

$$\|\mu w_{xx}\|_{L^2(0, T; H^{-1}(0, L))}^2 \leq \sup_{t \in [0, T]} |\mu|^2 \|w\|_{L^2(0, T; H^1(0, L))}^2.$$

ii) The case $s = 1$. Following the previous steps, we have

$$\begin{aligned} & \| (uw)_x \|_{L^2(Q)}^2 \\ & \leq 2 \int_0^T \left(\|w(\cdot, t)\|_{L^\infty(0, L)}^2 \|u(\cdot, t)\|_{H^1(0, L)}^2 + \|u(\cdot, t)\|_{L^\infty(0, L)}^2 \|w(\cdot, t)\|_{H^1(0, L)}^2 \right) dt \\ & \leq C \|u\|_{Y_T^1}^2 \|w\|_{Y_T^1}^2 \end{aligned}$$

and

$$\|\mu w_{xx}\|_{L^2(Q)}^2 \leq \sup_{t \in [0, T]} |\mu|^2 \|w\|_{L^2(0, T; H^2(0, L))}^2 \leq C \|\mu\|_{L^\infty(0, T)}^2 \|w\|_{Y_T^1}^2.$$

Similar arguments for $s = 2, 3$ as well as interpolation properties allow to complete the proof. $\square$

Next we consider the following linear problem

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} = f & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L), \end{cases} \quad (22)$$

where $\mu(t) \geq \mu_0 > 0$ for $t \in [0, T]$. For given $t \in [0, T]$, consider the operator $A(t) := -\partial_x^3 + \mu(t)\partial_x^2$ defined on

$$\mathcal{D}(A(t)) := \mathcal{D}(A) = \{u \in H^3(0, L) \cap H_0^1(0, L) : u(0) = u(L) = 0, u_x(0) = u_x(L)\} \subset L^2(0, L).$$

For any $\varphi \in \mathcal{D}(A)$,

$$\langle A(t)\varphi, \varphi \rangle_{L^2(0,L)} = - \int_0^L \varphi_{xxx} \varphi \, dx + \mu(t) \int_0^L \varphi_{xx} \varphi \, dx \leq -\mu(t) \int_0^L |\varphi_x|^2 \, dx \leq 0.$$

Thus the operator $A(t)$ is the infinitesimal generator of a C_0 semigroup of contractions in $L^2(0, L)$ and the family of the operators $\{A(t)\}_{0 \leq t \leq T}$ is stable (cf. Definition 2.1 on page 129 of [30]) and satisfies the assumptions of Theorem 3.1 in [30] on page 135. For any $y_0 \in L^2(0, L)$ and $f \in L^1(0, T; L^2(0, L))$, (22) admits a unique mild solution $y \in C([0, T]; L^2(0, L))$, given by

$$y(t) = U(t, 0)y_0 + \int_0^t U(t, \tau)f(\tau)d\tau \quad \forall t \in [0, T],$$

where $U(t, s)$, $0 \leq s \leq t \leq T$ is the associated evolution system. Using the same arguments as in the proofs of Lemma 2.1 and Proposition 2.2, we can show the following estimates hold for solutions of the system (22).

Lemma 2.7. *For every $T > 0$, $f \in L^1(0, T; L^2(0, L))$ and $y_0 \in L^2(0, L)$, the corresponding mild solution of (22) belongs to $C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))$ and satisfies*

$$\|y\|_{L^\infty(0,T;L^2(0,L))} + \|y\|_{L^2(0,T;H^1(0,L))} \leq C_1(\|y_0\|_{L^2(0,L)} + \|f\|_{L^1(0,T;L^2(0,L))})$$

for some constant $C_1 > 0$. Moreover, for given $0 \leq s \leq 3$, if $(f, y_0) \in L^2(0, T; H^{s-1}(0, L)) \times H^s(0, L)$, then

$$y \in C([0, T]; H^s(0, L)) \cap L^2(0, T; H^{s+1}(0, L))$$

and

$$\|y\|_{L^2(0,T;H^{s+1}(0,L))} \leq C_2 \left(\|f\|_{L^2(0,T;H^{s-1}(0,L))} + \|y_0\|_{H^s(0,L)} \right), \quad (23)$$

for some constant $C_2 > 0$.

Proposition 2.3. *Let $T > 0$ and $s \in [0, 3]$ be given and assume $\bar{y} \in Y_T^s$. Then for any $y_0 \in H^s(0, L)$, the linearized system (10) admits a unique solution $y \in Y_T^s$.*

Proof. The proof is developed for the case $s = 0$. Similar arguments allow to extend this result for $0 < s \leq 3$. Let us consider $R > 0$ and $0 < \theta \leq \min\{1, T\}$ two appropriate constants to be determined. Let $B_{\theta,R} := \{z \in Y_\theta^0 : \|z\|_{Y_\theta^0} \leq R\}$ and define a map $\Lambda : B_{\theta,R} \rightarrow B_{\theta,R}$ by $\Lambda(z) = y$, where y is the unique solution of

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} = -(\bar{y}z)_x & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L). \end{cases}$$

Obviously,

$$\Lambda(z) = U(t, 0)y_0 - \int_0^t U(t, \tau)((\bar{y}z)_x)(\tau) d\tau.$$

From the above representation and using Lemma 2.7, there exist positive constants C_3, C_4 such that

$$\|\Lambda(z)\|_{Y_\theta^0} \leq C_3 \|y_0\|_{L^2(0,L)} + C_4 \theta^{1/2} \|\bar{y}\|_{Y_T^0} \|z\|_{Y_\theta^0}. \quad (24)$$

Choose $R > 0$ and $T^* = \theta$ such that

$$R = 2C_3 \|y_0\|_{L^2(0,L)} \quad \text{and} \quad C_4 T^{*1/2} \|\bar{y}\|_{Y_T^0} \leq \frac{1}{4}.$$

Then, by (24) we have that $\|\Lambda(z)\|_{Y_{T^*}^0} \leq R$. Furthermore, for every $z^1, z^2 \in B_{T^*,R}$,

$$\|\Lambda(z^1) - \Lambda(z^2)\|_{Y_{T^*}^0} \leq 2C_4 T^{*1/2} \|\bar{y}\|_{Y_T^0} \|z^1 - z^2\|_{Y_T^0} \leq \frac{1}{2} \|z^1 - z^2\|_{Y_{T^*}^0}.$$

Therefore, Λ is a contraction mapping on $B_{T^*,R}$ and it has a unique fixed point $z^2 \in Y_{T^*}^0$ which is the solution to the linearized problem (2.2) in $(0, T^*)$. Finally, we can observe that $T^* \in (0, T)$ is independent on $\|y_0\|_{L^2(0,L)}$, it implies that the previous arguments can be extended on intervals $(T^*, 2T^*], (2T^*, 3T^*], \dots, ((n-1)T^*, nT^* = T]$. Therefore, the existence of a unique solution of (10) in $(0, T)$ is guaranteed. This completes the proof of Proposition 2.3. $\square$

Remark 2.2. As consequence of Proposition 2.3 and Lemma 2.7, for any trajectory $\bar{y} \in Y_T^s$, the solution y of (10) satisfies

$$\|y\|_{Y_T^s} \leq C \left(\|f\|_{L^2(0,T;H^{s-1}(0,L))} + \|y_0\|_{H^s(0,L)} \right), \quad (25)$$

for some positive constant C .

2.2. Nonlinear case

In this subsection we turn to consider the following nonlinear initial boundary value problem (IBVP):

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} + yy_x = 0 & \text{in } (0, L) \times (0, +\infty), \\ y(0, t) = y(L, t) = 0 & \text{in } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L). \end{cases} \quad (26)$$

Proposition 2.4. Let $s \in [0, 3]$ and $T > 0$ be given. There exists $\delta > 0$ such that for any $y_0 \in H^s(0, L)$ satisfying $\|y_0\|_{H^s(0,L)} \leq \delta$, the nonlinear system (26) admits a unique solution $y \in Y_T^s$.

Proof. The proof follows the same scheme of the linear case. In fact, let $R > 0$ be an appropriate constant to be determined. Again, we consider a map $\Lambda : B_R \subset Y_T^s \rightarrow Y_T^s$ by $\Lambda(z) = y$ where y solves

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} = -zz_x & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{in } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L). \end{cases}$$

In this case,

$$\Lambda(z) = U(t, 0)y_0 - \int_0^t U(t, \tau)(zz_x)(\tau) d\tau.$$

Using Lemma 2.7, Lemma 2.6 and (25), there exist positive constants C_5, C_6 such that

$$\|\Lambda(z)\|_{Y_T^s} \leq C_5 \|y_0\|_{H^s(0, L)} + C_6 \|z\|_{Y_T^s}^2. \quad (27)$$

Consider $R > 0$ such that

$$R = 2C_5 \|y_0\|_{H^s(0, L)} \quad \text{and} \quad C_6 R \leq \frac{1}{4}. \quad (28)$$

From (28), it is enough to define $\delta := (8C_5 C_6)^{-1}$. Then, by (27) we have that $\|\Lambda(z)\|_{Y_T^s} \leq R$. Furthermore, for every $z^1, z^2 \in B_R$,

$$\begin{aligned} \|\Lambda(z^1) - \Lambda(z^2)\|_{Y_T^s} &\leq C_6 \|z^1 z_x^1 - z^2 z_x^2\|_{L^2(0, T; H^{s-1}(0, L))} \\ &\leq C_6 (\|z^1\|_{Y_T^s} + \|z^2\|_{Y_T^s}) \|z^1 - z^2\|_{Y_T^s} \\ &\leq \frac{1}{2} \|z^1 - z^2\|_{Y_T^s}. \end{aligned}$$

Therefore Λ is a contraction mapping on B_R and it has a unique fixed point $z^2 \in Y_T^s$ which is the solution of (26). $\square$

3. Carleman inequality

In this section we will prove the Carleman estimate given in Theorem 1.1. To do this, we introduce weight functions defined as follows. Let ω be a nonempty open subset of $(0, L)$ and ϕ a positive function in $[0, L]$ such that $\phi \in C^4([0, L])$ and satisfies

$$\phi(0) = \phi(L), \quad \phi'(0) < 0, \quad \phi'(L) > 0, \quad |\phi'(0)| = |\phi'(L)|, \quad (29)$$

$$\phi'' < 0 \quad \text{in } \overline{(0, L) \setminus \omega}. \quad (30)$$

Thus, we consider

$$\begin{aligned}\alpha(x, t) &:= \phi(x)\xi(t), \quad \xi(t) := \frac{1}{t^2(T-t)^2}, \\ \hat{\alpha}(t) &:= \max_{x \in [0, L]} \alpha(x, t), \quad \check{\alpha}(t) := \min_{x \in [0, L]} \alpha(x, t), \quad 2\hat{\alpha}(t) < 3\check{\alpha}(t).\end{aligned}\tag{31}$$

For instance, if $\omega = (\ell_1, \ell_2) \subset (0, L)$, then it is easy to verify that ϕ defined as follows satisfies (29) and (30):

$$\phi(x) := \begin{cases} x^3 - 3\ell_1 x^2 - x + 2L^3 + L + d & \text{if } x \in [0, \ell_1], \\ -x^3 + (1 + 3L^2)x + d & \text{if } x \in [\ell_2, L]. \end{cases}$$

Additionally, if $d > \max\{4L^3 + 2L + 3\ell_1^3 + 3\ell_1, 4L^3 + 2L + 3\ell_2^3\}$, and

$$\begin{aligned}\max_{x \in [0, L]} \phi(x) &= \phi(0) = 2L^3 + L + d, \\ \min_{x \in [0, L]} \phi(x) &= \min\{\phi(\ell_1) - \ell_1^3 + 2L^3 - L, \phi(\ell_2) - (1 + 3L^2)\ell_2^2\},\end{aligned}$$

the inequality $2\hat{\alpha}(t) < 3\check{\alpha}(t)$ can be verified.

Proof of Theorem 1.1. For an easier comprehension, we divide the proof in several steps:

Step 1. Decomposition of the solution. In this step, we decompose the solution φ of (8) in order to obtain L^2 regularity on the right-hand side of (8). In other words, let us introduce z and u , the solutions of the following systems

$$\begin{cases} -z_t - z_{xxx} - \mu(t)z_{xx} - \bar{y}z_x = -(\rho_0)_t \varphi & \text{in } Q, \\ z(0, t) = z(L, t) = 0 & \text{on } (0, T), \\ z_x(0, t) = z_x(L, t) & \text{on } (0, T), \\ z(\cdot, T) = 0 & \text{in } (0, L), \end{cases}\tag{32}$$

and

$$\begin{cases} -u_t - u_{xxx} - \mu(t)u_{xx} - \bar{y}u_x = \rho_0 g & \text{in } Q, \\ u(0, t) = u(L, t) = 0 & \text{on } (0, T), \\ u_x(0, t) = u_x(L, t) & \text{on } (0, T), \\ u(\cdot, T) = 0 & \text{in } (0, L), \end{cases}\tag{33}$$

where $\rho_0(t) = e^{-s\hat{\alpha}}$. By uniqueness for the linear KdVB equation, we have

$$\rho_0 \varphi = u + z.\tag{34}$$

The rest of the proof consists in making a Carleman inequality for the system (32), meanwhile, for the system (33) we will use the regularity result (25), namely

$$\|u\|_{L^2(0, T; H^2(0, L))}^2 \leq C \|\rho_0 g\|_{L^2(Q)}^2.\tag{35}$$

Step 2. Change of variables and decomposition of a special operator. In this step, we consider the differential operator satisfied by a new variable w , which will be z up to a weight function. More precisely, let us define $w = e^{-s\alpha}z$. In this way, if L is the operator defined by $L := \partial_t + \partial_{xxx} + \mu(t)\partial_{xx}$, the identity $e^{-s\alpha}L(e^{s\alpha}w) = e^{-s\alpha}((\rho_0)_t\varphi - \bar{y}z_x)$ is equivalent to:

$$L_1w + L_2w + Rw = G,$$

where

$$\begin{aligned} L_1w &= w_t + w_{xxx} + 3s^2(\alpha_x)^2w_x, \\ L_2w &= 3s(\alpha_x)w_{xx} + s^3(\alpha_x)^3w + 3s(\alpha_{xx})w_x, \\ G &= e^{-s\alpha}((\rho_0)_t\varphi - \bar{y}z_x) \end{aligned} \quad (36)$$

and

$$Rw := \mu(t)s\alpha_{xx}w + s\alpha_t w + s\alpha_{xxx}w + 3s^2\alpha_x\alpha_{xx}w + \mu(t)(2s\alpha_xw_x + w_{xx} + s^2\alpha_x^2w).$$

Therefore,

$$\|L_1w\|_{L^2(Q)}^2 + \|L_2w\|_{L^2(Q)}^2 + 2\langle L_1w, L_2w \rangle = \|G - Rw\|_{L^2(Q)}^2, \quad (37)$$

where $\langle \cdot, \cdot \rangle$ is the $L^2(Q)$ inner product. In the next step, we will estimate the terms that arise of the inner product $\langle L_1w, L_2w \rangle$. This will give an inequality with global terms on the left-hand side, meanwhile the local terms will appear on the right-hand side. Finally, after returning to the principal variable z , the local terms will be estimated using bootstrap arguments based on the smoothing of the KdVB equation.

Step 3. First estimates. In this step, we develop the nine terms appearing in $\langle L_1w, L_2w \rangle$. Using integration by parts, we have:

$$\begin{aligned} I^{1,1} &:= \langle L_1^1w, L_2^1w \rangle = 3s \iint_Q \alpha_x w_t w_{xx} dx dt \\ &= -3s \iint_Q \alpha_{xx} w_t w_x dx dt + \frac{3s}{2} \iint_Q \alpha_{xt} |w_x|^2 dx dt. \\ I^{1,2} &:= \langle L_1^1w, L_2^2w \rangle = s^3 \iint_Q (\alpha_x)^3 w_t w dx dt = -\frac{3s^3}{2} \iint_Q (\alpha_x)^2 \alpha_{xt} |w|^2 dx dt. \\ I^{1,3} &:= \langle L_1^1w, L_2^3w \rangle = 3s \iint_Q \alpha_{xx} w_x w_t dx dt \\ &= -3s \iint_Q \alpha_x w_x w_{xt} dx dt - 3s \iint_Q \alpha_x w_{xx} w_t dx dt \\ &= \frac{3s}{2} \iint_Q \alpha_{xt} |w_x|^2 dx dt - I^{1,1}. \end{aligned} \quad (38)$$

$$\begin{aligned}
 I^{2,1} &:= \langle L_1^2 w, L_2^1 w \rangle = 3s \iint_Q \alpha_x w_{xx} w_{xxx} dx dt \\
 &= -\frac{3s}{2} \iint_Q \alpha_{xx} |w_{xx}|^2 dx dt + \underbrace{\frac{3s}{2} \int_0^T \left(\alpha_x |w_{xx}|^2 \Big|_{x=0}^{x=L} \right) dt}_A.
 \end{aligned} \tag{39}$$

$$\begin{aligned}
 I^{2,2} &:= \langle L_1^2 w, L_2^2 w \rangle = s^3 \iint_Q (\alpha_x)^3 w w_{xxx} dx dt \\
 &= -3s^3 \iint_Q (\alpha_x)^2 \alpha_{xx} w w_{xx} dx dt - s^3 \iint_Q (\alpha_x)^3 w_x w_{xx} dx dt \\
 &= 3s^3 \iint_Q ((\alpha_x)^2 \alpha_{xx})_x w w_x dx dt + 3s^3 \iint_Q (\alpha_x)^2 \alpha_{xx} |w_x|^2 dx dt \\
 &\quad + \frac{s^3}{2} \iint_Q ((\alpha_x)^3)_x |w_x|^2 dx dt - \frac{s^3}{2} \int_0^T \left((\alpha_x)^3 |w_x|^2 \Big|_{x=0}^{x=L} \right) dt \\
 &= -\frac{3s^3}{2} \iint_Q ((\alpha_x)^2 \alpha_{xx})_{xx} |w|^2 dx dt \\
 &\quad + \frac{9s^3}{2} \iint_Q (\alpha_x)^2 \alpha_{xx} |w_x|^2 dx dt - \underbrace{\frac{s^3}{2} \int_0^T \left((\alpha_x)^3 |w_x|^2 \Big|_{x=0}^{x=L} \right) dt}_B,
 \end{aligned} \tag{40}$$

$$\begin{aligned}
 I^{2,3} &:= \langle L_1^2 w, L_2^3 w \rangle = 3s \iint_Q \alpha_{xx} w_x w_{xxx} dx dt \\
 &= -3s \iint_Q \alpha_{xxx} w_x w_{xx} dx dt - 3s \iint_Q \alpha_{xx} |w_{xx}|^2 dx dt + 3s \int_0^T \left(\alpha_{xx} w_x w_{xx} \Big|_{x=0}^{x=L} \right) dt \\
 &= -3s \iint_Q \alpha_{xx} |w_{xx}|^2 dx dt + \frac{3s}{2} \iint_Q \alpha_{xxx} |w_x|^2 dx dt + \tilde{C},
 \end{aligned} \tag{41}$$

where

$$\tilde{C} := 3s \int_0^T \left(\alpha_{xx} w_x w_{xx} \Big|_{x=0}^{x=L} \right) dt - \frac{3s}{2} \int_0^T \left(\alpha_{xxx} |w_x|^2 \Big|_{x=0}^{x=L} \right) dt.$$

$$\begin{aligned}
I^{3,1} &:= \langle L_1^3 w, L_2^1 w \rangle = 9s^3 \iint_Q (\alpha_x)^3 w_x w_{xx} dx dt \\
&= -\frac{9s^3}{2} \iint_Q ((\alpha_x)^3)_x |w_x|^2 dx dt + \frac{9s^3}{2} \int_0^T \left((\alpha_x)^3 |w_x|^2 \Big|_{x=0}^{x=L} \right) dt \\
&= -\frac{27s^3}{2} \iint_Q (\alpha_x)^2 \alpha_{xx} |w_x|^2 dx dt + \underbrace{\frac{9s^3}{2} \int_0^T \left((\alpha_x)^3 |w_x|^2 \Big|_{x=0}^{x=L} \right) dt}_D.
\end{aligned} \tag{42}$$

$$\begin{aligned}
I^{3,2} &:= \langle L_1^3 w, L_2^2 w \rangle = 3s^5 \iint_Q (\alpha_x)^5 w w_x dx dt \\
&= -\frac{3s^5}{2} \iint_Q ((\alpha_x)^5)_x |w|^2 dx dt \\
&= -\frac{15s^5}{2} \iint_Q (\alpha_x)^4 \alpha_{xx} |w|^2 dx dt.
\end{aligned} \tag{43}$$

$$I^{3,3} := \langle L_1^3 w, L_2^3 w \rangle = 9s^3 \iint_Q (\alpha_x)^2 \alpha_{xx} |w_x|^2 dx dt. \tag{44}$$

Putting together all the computations, we have

$$\begin{aligned}
\langle L_1 w, L_2 w \rangle &= -\frac{15s^5}{2} \iint_Q (\alpha_x)^4 \alpha_{xx} |w|^2 dx dt \\
&\quad - \frac{3s^3}{2} \iint_Q ((\alpha_x)^2 \alpha_{xx})_{xx} |w|^2 dx dt - \frac{3s^3}{2} \iint_Q (\alpha_x)^2 \alpha_{xt} |w|^2 dx dt \\
&\quad + \frac{3s}{2} \iint_Q \alpha_{xxx} |w_x|^2 dx dt + \frac{3s}{2} \iint_Q \alpha_{xt} |w_x|^2 dx dt \\
&\quad - \frac{9s}{2} \iint_Q \alpha_{xx} |w_{xx}|^2 dx dt + A + B + \tilde{C} + D.
\end{aligned} \tag{45}$$

Now, taking into account the first boundary condition of (32) and (30), we can deduce the inequalities:

$$Cs^5 \iint_Q \xi^5 |w|^2 dx dt - Cs^5 \iint_{\omega \times (0,T)} \xi^5 |w|^2 dx dt \leq -\frac{15s^5}{2} \iint_Q (\alpha_x)^4 \alpha_{xx} |w|^2 dx dt \tag{46}$$

and

$$-\frac{9s}{2} \iint_Q \alpha_{xx} |w_{xx}|^2 dx dt \geq Cs \iint_Q \xi |w_{xx}|^2 dx dt - Cs \iint_{\omega \times (0, T)} \xi |w_{xx}|^2 dx dt, \quad (47)$$

for any $s \geq C(L, \omega, T)$.

On the other hand, integrating by parts and using Young's inequality, we also obtain

$$s^3 \iint_Q \xi^3 |w_x|^2 dx dt \leq \iint_Q (s^5 \xi^5 |w|^2 + s \xi |w_{xx}|^2) dx dt. \quad (48)$$

Hence, combining (37) and (45)–(48), we have that there exists $C = C(L, \omega, T)$ such that

$$\begin{aligned} & \|L_1 w\|_{L^2(Q)}^2 + \|L_2 w\|_{L^2(Q)}^2 + \iint_Q (s^5 \xi^5 |w|^2 + s^3 \xi^3 |w_x|^2 + s \xi |w_{xx}|^2) dx dt + A + B + \tilde{C} + D \\ & \leq C \left(\iint_{\omega \times (0, T)} ((s \xi)^5 |w|^2 + s \xi |w_{xx}|^2) dx dt + \|G\|_{L^2(Q)}^2 + \|Rw\|_{L^2(Q)}^2 \right), \end{aligned} \quad (49)$$

for any $s \geq C(L, \omega, T)$.

Note that the last term on the right-hand side (49) can be absorbed by the left-hand side for $s \geq C(L, \omega, T, \|\mu\|_{L^\infty(0, T)})$. Furthermore, since $G = [(\rho_0)_t \varphi - \bar{y} z_x] e^{-s\alpha}$, we can estimate the term $\bar{y} z_x$ using the identity $w_x + s\alpha_x w = e^{-s\alpha} z_x$ and the inequality

$$|\bar{y} z_x|^2 e^{-2s\alpha} \leq Cs |\bar{y} w_x|^2 + Cs^2 (\alpha_x)^2 |\bar{y} w|^2.$$

Additionally, from (34) and (35), we also have that $|(\rho_0)_t \varphi| \leq Cs \xi^{3/2} |\rho_0 \varphi|$. Thus, we readily have that there exists constant $C = C(L, \omega, T, \|\mu\|_{L^\infty(0, T)}, \|\bar{y}\|_{C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))})$ such that

$$\begin{aligned} & \iint_Q (s^5 \xi^5 |w|^2 + s^3 \xi^3 |w_x|^2 + s \xi |w_{xx}|^2) dx dt + A + B + \tilde{C} + D \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\alpha}} dx dt + \iint_{\omega \times (0, T)} ((s \xi)^5 |w|^2 + s \xi |w_{xx}|^2) dx dt \right). \end{aligned}$$

Finally, using the weight functions defined in (29) and (30), the terms $A - D$ in the previous inequality can be estimated as follows:

$$\begin{aligned} A &= \frac{3s}{2} \int_0^T (\alpha_x(L, t) |w_{xx}(L, t)|^2 - \alpha_x(0, t) |w_{xx}(0, t)|^2) dt \\ &\geq Cs \int_0^T \xi (|w_{xx}(0, t)|^2 + |w_{xx}(L, t)|^2) dt. \end{aligned}$$

$$\begin{aligned} B + D &= 4s^3 \int_0^T ((\alpha_x(L, t))^3 |w_x(L, t)|^2 - (\alpha_x(0, t))^3 |w_x(0, t)|^2) dt \\ &\geq Cs^3 \int_0^T \xi^3 |w_x(L, t)|^2 dt, \end{aligned}$$

and

$$\begin{aligned} \tilde{C} &= 3s \int_0^T (\alpha_{xx}(L, t) w_x(L, t) w_{xx}(L, t) - \alpha_{xx}(0, t) w_x(0, t) w_{xx}(0, t)) dt \\ &\quad - \frac{3s}{2} \int_0^T (\alpha_{xxx}(L, t) |w_x(L, t)|^2 - \alpha_{xxx}(0, t) |w_x(0, t)|^2) dt \\ &\leq Cs^2 \int_0^T \xi |w_x(L, t)|^2 dt + C \int_0^T \xi (|w_{xx}(0, t)|^2 + |w_{xx}(L, t)|^2) dt. \end{aligned}$$

Therefore, at this moment we have the following inequality

$$\begin{aligned} &\iint_Q (s^5 \xi^5 |w|^2 + s^3 \xi^3 |w_x|^2 + s \xi |w_{xx}|^2) dx dt + s^3 \int_0^T \xi^3 |w_x(L, t)|^2 dt \\ &\quad + s \int_0^T \xi (|w_{xx}(0, t)|^2 + |w_{xx}(L, t)|^2) dt \\ &\leq C \left(\iint_Q |g|^2 e^{-2s\hat{a}} dx dt + \iint_{\omega \times (0, T)} (s^5 \xi^5 |w|^2 + s \xi |w_{xx}|^2) dx dt \right), \end{aligned} \quad (50)$$

for any $s \geq C(L, \omega, T, \|\mu\|_{L^\infty(0, T)}, \|\bar{\gamma}\|_{C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))})$.

Step 4. Local estimates. In this step, we turn back to our original function and use bootstrap arguments as in [7] and [21] to estimate the local term associated to $|w_{xx}|$.

Recall that $z = e^{s\alpha} w$. Then, a direct computation allow to obtain

$$|z_x|^2 e^{-2s\alpha} \leq C(s^2 \xi^2 |w|^2 + |w_x|^2) \quad (51)$$

and

$$|z_{xx}|^2 e^{-2s\alpha} \leq C(s^4 \xi^4 |w|^2 + s^2 \xi^2 |w_x|^2 + |w_{xx}|^2). \quad (52)$$

On the other hand,

$$|w_{xx}|^2 \leq C e^{-2s\alpha} (s^4 \xi^4 |z|^2 + s^2 \xi^2 |z_x|^2 + |z_{xx}|^2). \quad (53)$$

From (53) the local term given in (50) can be written by

$$\iint_{\omega \times (0, T)} (s^5 \xi^5 |z|^2 + s^3 \xi^3 |z_x|^2 + s \xi |z_{xx}|^2) e^{-2s\alpha} dx dt. \quad (54)$$

In addition, the weight functions $\hat{\alpha}$, $\check{\alpha}$, ξ and (50)–(54) allow us to deduce the following inequality

$$\begin{aligned} & \iint_Q (s^5 \xi^5 |z|^2 + s^3 \xi^3 |z_x|^2 + s \xi |z_{xx}|^2) e^{-2s\hat{\alpha}} dx dt \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\alpha}} dx dt + \iint_{\omega \times (0, T)} (s^5 \xi^5 |z|^2 + s^3 \xi^3 |z_x|^2 + s \xi |z_{xx}|^2) e^{-2s\check{\alpha}} dx dt \right), \end{aligned} \quad (55)$$

for any $s \geq C(L, \omega, T, \|\mu\|_{L^\infty(0, T)}, \|\bar{\gamma}\|_{C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))})$.

Using that $H^1(\omega) = (H^3(\omega), L^2(\omega))_{2/3, 2}$ and $H^2(\omega) = (H^3(\omega), L^2(\omega))_{1/3, 2}$, the last two terms in the right-hand side of (55) can be upper bounded as follows:

$$s^3 \iint_{\omega \times (0, T)} \xi^3 e^{-2s\check{\alpha}} |z_x|^2 dx dt \leq \underbrace{s^3 \int_0^T \xi^3 e^{-2s\check{\alpha}} \|z\|_{L^2(\omega)}^{4/3} \|z\|_{H^3(\omega)}^{2/3} dt}_{J_1}$$

and

$$s \iint_{\omega \times (0, T)} \xi e^{-2s\check{\alpha}} |z_{xx}|^2 dx dt \leq \underbrace{s \int_0^T \xi e^{-2s\check{\alpha}} \|z\|_{L^2(\omega)}^{2/3} \|z\|_{H^3(\omega)}^{4/3} dt}_{J_2}.$$

Now, applying Young's inequality

$$J_1 \leq C(\varepsilon) s^{11/2} \int_0^T \xi^{11/2} e^{-3s\check{\alpha} + s\hat{\alpha}} \|z\|_{L^2(\omega)}^2 dt + \varepsilon s^{-2} \int_0^T \xi^{-2} e^{-2s\hat{\alpha}} \|z\|_{H^3(\omega)}^2 dt$$

and

$$J_2 \leq C(\varepsilon) s^7 \int_0^T \xi^7 e^{-6s\check{\alpha} + 4s\hat{\alpha}} \|z\|_{L^2(\omega)}^2 dt + \varepsilon s^{-2} \int_0^T \xi^{-2} e^{-2s\hat{\alpha}} \|z\|_{H^3(\omega)}^2 dt,$$

for any $\varepsilon > 0$.

Putting together (55) and the previous estimates, we have

$$\begin{aligned} & \iint_Q (s^5 \xi^5 |z|^2 + s^3 \xi^3 |z_x|^2 + s \xi |z_{xx}|^2) e^{-2s\hat{\alpha}} dx dt \\ & \leq C \iint_Q |g|^2 e^{-2s\hat{\alpha}} dx dt + C s^7 \iint_{\omega \times (0, T)} \xi^7 e^{-6s\hat{\alpha} + 4s\hat{\alpha}} |z|^2 dx dt + \varepsilon \left(s^{-2} \int_0^T \xi^{-2} e^{-2s\hat{\alpha}} \|z\|_{H^3(\omega)}^2 dt \right). \end{aligned} \quad (56)$$

Finally, in order to estimate the associated terms to $\|z\|_{H^3(\omega)}^2$, we will use a bootstrap argument based on the smoothing effect of the KdVB equation. Let us start by defining $\tilde{z} := \tilde{\rho}(t)z$ with $\tilde{\rho}(t) := s^{1/2} \xi e^{-s\hat{\alpha}}$. From (32), we see that $\tilde{z}$ is the solution of the system

$$\begin{cases} -\tilde{z}_t - \tilde{z}_{xxx} - \mu(t)\tilde{z}_{xx} - \bar{y}\tilde{z}_x = \tilde{\rho}(\rho_0)_t \varphi - \tilde{\rho}_t z & \text{in } Q, \\ \tilde{z}(0, t) = \tilde{z}(L, t) = 0 & \text{on } (0, T), \\ \tilde{z}_x(0, t) = \tilde{z}_x(L, t) & \text{on } (0, T), \\ \tilde{z}(\cdot, T) = 0 & \text{in } (0, L). \end{cases} \quad (57)$$

Taking into account the estimates $|\tilde{\rho}_t| \leq C s^{3/2} \xi^{5/2} e^{-s\hat{\alpha}}$, $|(\rho_0)_t| \leq C s \xi^{3/2} e^{-s\hat{\alpha}}$, and the regularity result (25), we can deduce that

$$\|\tilde{z}\|_{L^2(0, T; H^2(0, L))}^2 \leq C \left(\|s^{3/2} \xi^{5/2} e^{-s\hat{\alpha}} z\|_{L^2(Q)}^2 + \|s^{3/2} \xi^{5/2} e^{-2s\hat{\alpha}} \varphi\|_{L^2(Q)}^2 \right). \quad (58)$$

The fact that $s^{3/2} \xi^{5/2} e^{-s\hat{\alpha}}$ is bounded allows us to use (35) and conclude that $\|\tilde{z}\|_{L^2(0, T; H^2(0, L))}^2$ is bounded by the left-hand side of (56) and $\|\rho_0 g\|_{L^2(Q)}^2$. Now, we define

$$\hat{z} := \hat{\rho}(t)z \quad \text{with} \quad \hat{\rho}(t) := s^{-1/2} \xi^{-1/2} e^{-s\hat{\alpha}}.$$

It is easy to see that $\hat{z}$ is the solution of (57) with $\tilde{\rho}$ replaced by $\hat{\rho}$. Besides, from (25) we get

$$\|\hat{z}\|_{L^2(0, T; H^3(0, L))}^2 \leq C \left(\|s^{1/2} \xi e^{-s\hat{\alpha}} z\|_{L^2((0, T); H^1(0, L))}^2 + \|s^{1/2} \xi e^{-2s\hat{\alpha}} \varphi\|_{L^2(0, T; H^1(0, L))}^2 \right). \quad (59)$$

Arguing as before, $\|\hat{z}\|_{L^2(0, T; H^3(0, L))}^2$ is bounded by the left-hand side of (56) and $\|\rho_0 g\|_{L^2(Q)}^2$.

By combining (56), (58) and (59), we obtain in particular

$$\begin{aligned} & \iint_Q (s^5 \xi^5 |z|^2 + s^3 \xi^3 |z_x|^2 + s \xi |z_{xx}|^2) e^{-2s\hat{\alpha}} dx dt + \|s^{-1/2} \xi^{-1/2} e^{-s\hat{\alpha}} z\|_{L^2(0, T; H^3(0, L))}^2 \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\alpha}} dx dt + s^7 \iint_{\omega \times (0, T)} \xi^7 e^{-6s\hat{\alpha} + 4s\hat{\alpha}} |z|^2 dx dt \right) + \varepsilon \left(s^{-2} \int_0^T \xi^{-2} e^{-2s\hat{\alpha}} \|z\|_{H^3(\omega)}^2 dt \right), \end{aligned} \quad (60)$$

for any $\varepsilon > 0$. For ε small enough, the last two terms in the right-hand side of (60) can be absorbed by the left-hand side. By returning to the variable φ the proof of Theorem 1.1 is ended. $\square$

4. Null controllability of the linearized system

In this section we will prove the null controllability for the system (7) with a right-hand side which decays exponentially to zero when t goes to T [16]. In other words, we would like to find $v \in L^2(0, T; L^2(\Omega))$ such that the solution of

$$\begin{cases} y_t + y_{xxx} - \mu(t)y_{xx} + \bar{y}y_x + y\bar{y}_x = h + v1_{\omega \times (0, T)} & \text{in } Q, \\ y(0, t) = y(L, t) = 0 & \text{on } (0, T), \\ y_x(0, t) = y_x(L, t) & \text{on } (0, T), \\ y(\cdot, 0) = y_0(\cdot) & \text{in } (0, L), \end{cases} \quad (61)$$

satisfies

$$y(\cdot, T) = 0 \quad \text{in } (0, L), \quad (62)$$

where the function h is in an appropriate weighted space. Before proving these results, we establish a Carleman inequality with weight functions not vanishing in $t = 0$. To do this, let $\ell(t) \in C^1([0, T])$ be a positive function in $[0, T)$ such that $\ell(t) = T^2/4$ for all $t \in [0, T/4]$ and $\ell(t) = t(T - t)$ for all $t \in [T/2, T]$. We introduce the following weight functions:

$$\begin{aligned} \beta(x, t) &= \phi(x)\tau(t), \quad \tau(t) = \frac{1}{\ell^2(t)}, \\ \hat{\beta}(t) &= \max_{x \in [0, L]} \beta(x, t), \quad \check{\beta}(t) = \min_{x \in [0, L]} \beta(x, t). \end{aligned} \quad (63)$$

Lemma 4.1. *There exist positive constants s, C with C depending on $s, \|\mu\|_{L^\infty(0, T)}, \omega, T$ such that every solution of (8) verifies*

$$\begin{aligned} & \iint_Q (\tau^5 |\varphi|^2 + \tau^3 |\varphi_x|^2 + \tau |\varphi_{xx}|^2) e^{-4s\hat{\beta}} dx dt + \|\varphi(0)\|_{L^2(0, L)}^2 \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\beta}} dx dt + \iint_{\omega \times (0, T)} \tau^7 e^{-6s\check{\beta} + 2s\hat{\beta}} |\varphi|^2 dx dt \right). \end{aligned} \quad (64)$$

Proof. By construction $\alpha = \beta$ and $\tau = \xi$ in $[0, L] \times (T/2, T)$, so that

$$\int_{T/2}^T \int_0^L (\xi^5 |\varphi|^2 + \xi^3 |\varphi_x|^2 + \xi |\varphi_{xx}|^2) e^{-4s\hat{\alpha}} dx dt = \int_{T/2}^T \int_0^L (\tau^5 |\varphi|^2 + \tau^3 |\varphi_x|^2 + \tau |\varphi_{xx}|^2) e^{-4s\hat{\beta}} dx dt.$$

As consequence of Theorem 1.1 we have the estimate

$$\begin{aligned} & \int_{T/2}^T \int_0^L (\tau^5 |\varphi|^2 + \tau^3 |\varphi_x|^2 + \tau |\varphi_{xx}|^2) e^{-4s\hat{\beta}} dx dt \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\alpha}} dx dt + \iint_{\omega \times (0, T)} \xi^7 e^{-6s\check{\alpha} + 2s\hat{\alpha}} |\varphi|^2 dx dt \right). \end{aligned}$$

Next, using that $\ell(t) = t(T - t)$ for any $t \in [T/2, T]$ and

$$e^{-2s\hat{\beta}} \geq C \quad \text{and} \quad \tau^7 e^{-6s\check{\beta} + 2s\hat{\beta}} \geq C \quad \text{in } [0, T/2],$$

we readily have

$$\begin{aligned} & \int_{T/2}^T \int_0^L (\tau^5 |\varphi|^2 + \tau^3 |\varphi_x|^2 + \tau |\varphi_{xx}|^2) e^{-2s\hat{\beta}} dx dt \\ & \leq C \left(\iint_Q |g|^2 e^{-2s\hat{\beta}} dx dt + \iint_{\omega \times (0, T)} \tau^7 e^{-6s\check{\beta} + 2s\hat{\beta}} |\varphi|^2 dx dt \right). \end{aligned} \quad (65)$$

On the other hand, by considering a function $\eta \in C^1([0, T])$ such that $\eta \equiv 1$ in $[0, T/2]$ and $\eta \equiv 0$ in $[3T/4, T]$, we can prove that $\eta\varphi$ satisfies the system

$$\begin{cases} -(\eta\varphi)_t - \eta\varphi_{xxx} - \mu(t)\eta\varphi_{xx} - \bar{y}\eta\varphi_x = -\eta g - \eta'\varphi & \text{in } Q, \\ (\eta\varphi)(0, t) = (\eta\varphi)(L, t) = 0 & \text{on } (0, T), \\ (\eta\varphi)_x(0, t) = (\eta\varphi)_x(L, t) & \text{on } (0, T), \\ (\eta\varphi)(\cdot, T) = 0 & \text{in } (0, L). \end{cases} \quad (66)$$

Additionally, from classical energy estimates and regularity result with right-hand side in $L^2(Q)$ (see (25)), we get

$$\|\varphi(0)\|_{L^2(0, L)}^2 + \|\varphi\|_{L^2(0, T/2; L^2(0, L))}^2 \leq C \left(\|g\|_{L^2(0, 3T/4; L^2(0, L))}^2 + \|\varphi\|_{L^2(T/2, 3T/4; L^2((0, L)))}^2 \right).$$

Taking into account that

$$\tau^5 e^{-2s\hat{\beta}} \geq C > 0, \quad \forall t \in [T/2, 3T/4] \quad \text{and} \quad e^{-4s\hat{\beta}} \geq C > 0, \quad \forall t \in [0, 3T/4],$$

we have

$$\begin{aligned} & \|\varphi(0)\|_{L^2(0, L)}^2 + \int_0^{T/2} \int_0^L (\tau^5 |\varphi|^2 + \tau^3 |\varphi_x|^2 + \tau |\varphi_{xx}|^2) e^{-4s\hat{\beta}} dx dt \\ & \leq C \left(\int_0^{3T/4} \int_0^L |g|^2 e^{-2s\hat{\beta}} dx dt + \int_{T/2}^{3T/4} \int_0^L \tau^5 e^{-4s\hat{\beta}} |\varphi|^2 dx dt \right). \end{aligned} \quad (67)$$

Putting together (65) and (67) we obtain the desired inequality (64). $\square$

Now, we can prove the null controllability of system (61). The idea is to look a solution y in a suitable weight functional space. To this end, we introduce the following space:

$$E := \{(y, v) : e^{s\hat{\beta}} y \in L^2(Q), \tau^{-7/2} e^{3s\check{\beta}-s\hat{\beta}} v 1_\omega \in L^2(Q), \\ e^{s\hat{\beta}} \tau^{-3/2} y \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L)), \\ e^{2s\hat{\beta}} \tau^{-5/2} (y_t + y_{xxx} - \mu(t)y_{xx} + \bar{y}y_x + y\bar{y}_x - v 1_\omega) \in L^2(0, T; H^{-1}(0, L))\}.$$

Proposition 4.1. *Consider $y_0 \in L^2(0, L)$ and $e^{2s\hat{\beta}} \tau^{-5/2} h \in L^2(Q)$. Then, there exists a function $v \in L^2(0, T; L^2(\omega))$ such that the associated solution (y, v) to (61) satisfies $(y, v) \in E$.*

Furthermore, there exists a positive constant C such that

$$\|v\|_{L^2(0,T;L^2(\omega))} \leq C(\|y_0\|_{L^2(0,L)} + \|h\|_{L^2(Q)}). \quad (68)$$

Proof. The proof follows some ideas [21] and therefore we only give a sketch of the proof. Let us now set

$$P_0 = \{\varphi \in C^3(\bar{Q}) : \varphi(0, t) = \varphi(L, t) = 0, \varphi_x(0, t) = \varphi_x(L, t), \text{ on } (0, T)\}$$

as well as the bilinear form

$$a(\hat{\varphi}, w) := \iint_Q e^{-2s\hat{\beta}} (L^* \hat{\varphi})(L^* w) dx dt + \iint_{\omega \times (0, T)} e^{-6s\check{\beta}+2s\hat{\beta}} \tau^7 \hat{\varphi} w dx dt, \quad \forall w \in P_0$$

and the linear form

$$\langle G, w \rangle := \iint_Q h w dx dt + \int_0^L y_0(\cdot) w(\cdot, 0) dx, \quad (69)$$

where L^* is the adjoint operator of L , i.e.,

$$L^* w = -w_t - w_{xxx} - \mu w_{xx} - \bar{w} w_x.$$

Note that Carleman inequality (64) holds for every $w \in P_0$, so that we have

$$\iint_Q \tau^5 e^{-4s\hat{\beta}} |w|^2 dx dt \leq C a(w, w), \quad \forall w \in P_0.$$

In consequence, it is very easy to prove that $a(\cdot, \cdot) : P_0 \times P_0 \rightarrow \mathbb{R}$ is a symmetric, definite positive bilinear form on P_0 , so that, by defining P as the completion of P_0 for the form induced by $a(\cdot, \cdot)$, it implies that $a(\cdot, \cdot)$ is well-defined, continuous and again definite positive on P . In addition, from Carleman inequality (64) and the hypothesis over the function h , i.e., $e^{2s\hat{\beta}} \tau^{-5/2} h \in L^2(Q)$, the linear form $w \rightarrow \langle G, w \rangle$ is well defined and continuous on P . Hence, Lax–Milgram’s lemma allows us to guarantee the existence and uniqueness of $\hat{\varphi} \in P$ satisfying

$$a(\hat{\varphi}, w) = \langle G, w \rangle; \forall w \in P. \quad (70)$$

Let us set

$$\begin{cases} \hat{y} &:= e^{-2s\hat{\beta}} L^* \hat{\varphi} & \text{in } Q, \\ \hat{v} &:= -e^{-6s\check{\beta}+2s\hat{\beta}} \tau^7 \hat{\varphi} & \text{in } \omega \times (0, T). \end{cases} \quad (71)$$

Observe that $\hat{y}$ verifies

$$a(\hat{\varphi}, \hat{\varphi}) = \iint_Q e^{2s\hat{\beta}} |\hat{y}|^2 dx dt + \iint_{\omega \times (0, T)} e^{6s\check{\beta}-2s\hat{\beta}} \tau^{-7} |\hat{v}|^2 dx dt < +\infty. \quad (72)$$

On the other hand, if v is replaced by $\hat{v}$ in (61), we can introduce $\tilde{y}$ as the weak solution of (61). It implies that $\tilde{y}$ is the unique solution of (61) with $v = \hat{v}$ defined by transposition (see Definition 2.1). Then $\tilde{y} = \hat{y}$ is the weak solution to (61).

Finally, we must verify that $(\hat{y}, \hat{v}) \in E$. Observe that, from (72) we have that $e^{s\hat{\beta}} \hat{y} \in L^2(Q)$ and $\tau^{-7/2} e^{3s\check{\beta}-s\hat{\beta}} \hat{v} \in L^2(Q)$. Moreover, the second hypothesis of Proposition 4.1 guarantees that

$$e^{2s\hat{\beta}} \tau^{-5/2} (\hat{y}_t + \hat{y}_{xxx} - \mu(t) \hat{y}_{xx} + \bar{y} \hat{y}_x + \hat{y} \bar{y}_x - \hat{v}) \in L^2(Q).$$

Thus, we must just check that $e^{s\hat{\beta}} \tau^{-3/2} \hat{y} \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))$. To do this, we define the functions

$$y^* := e^{s\hat{\beta}} \tau^{-3/2} \hat{y} \quad \text{and} \quad h^* := e^{s\hat{\beta}} \tau^{-3/2} (h + \hat{v}).$$

Observe that y^* satisfies the system

$$\begin{cases} y_t^* + y_{xxx}^* - \mu(t) y_{xx}^* + \bar{y} y_x^* + y^* \bar{y}_x = h^* + (e^{s\hat{\beta}} \tau^{-3/2})_t \hat{y} & \text{in } Q, \\ y^*(0, t) = y^*(L, t) = 0 & \text{on } (0, T), \\ y_x^*(0, t) = y_x^*(L, t) & \text{on } (0, T), \\ y^*(\cdot, 0) = e^{s\hat{\beta}(0)} \tau^{-3/2}(0) \hat{y}_0(\cdot) & \text{in } (0, L). \end{cases}$$

Since $e^{s\hat{\beta}} h \in L^2(Q)$ and $2\hat{\beta} < 3\check{\beta}$ (see (31)), we obtain that $h^* + (e^{s\hat{\beta}} \tau^{-3/2})_t \hat{y} \in L^2(Q)$, in particular in $L^2(0, T; H^{-1}(0, L))$. Furthermore, for $\hat{y}_0 \in L^2(0, L)$, Proposition 2.3 allows us to have $y^* \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))$.

By considering $\hat{v}$ defined in (71), the bilinear form (69) and the identity (70), we can deduce (68). This concludes the sketch of the proof of Proposition 4.1. $\square$

5. Local exact controllability to trajectories

In this section we give the proof of Theorem 1.2 through fixed point arguments. In order to apply the results obtained in the previous sections we consider the following change of variable. Let us set $y - \bar{y} =: z$ and use this equality in (2), where $\bar{y}$ solves (3). It is easy to verify that z satisfies

$$\begin{cases} z_t + z_{xxx} - \mu(t)z_{xx} + (z\bar{y})_x + zz_x = v1_\omega & \text{in } Q, \\ z(0, t) = z(L, t) = 0 & \text{on } (0, T), \\ z_x(0, t) = z_x(L, t) & \text{on } (0, T), \\ z(\cdot, 0) = y_0 - \bar{y}_0 & \text{in } (0, L). \end{cases} \quad (73)$$

Observe that these changes reduce our problem to a local null controllability for the solution z of the nonlinear problem (73), i.e., we are looking a function control v such that z solution of (73) satisfies

$$z(\cdot, T) = 0 \quad \text{in } (0, L). \quad (74)$$

To do this, we will use the following inverse mapping theorem (see [1]).

Theorem 5.1. *Suppose that $\mathcal{B}_1, \mathcal{B}_2$ are Banach spaces and $\mathcal{A} : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ is a continuously differentiable map. We assume that for $b_1^0 \in \mathcal{B}_1, b_2^0 \in \mathcal{B}_2$ the equality*

$$\mathcal{A}(b_1^0) = b_2^0 \quad (75)$$

holds and $\mathcal{A}'(b_1^0) : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ is an epimorphism. Then there exists $\delta > 0$ such that for any $b_2 \in \mathcal{B}_2$ which satisfies the condition $\|b_2^0 - b_2\|_{\mathcal{B}_2} < \delta$ there exists a solution $b_1 \in \mathcal{B}_1$ of the equation

$$\mathcal{A}(b_1) = b_2.$$

In our framework, we use the above theorem with the spaces

$$\mathcal{B}_1 := E \quad \text{and} \quad \mathcal{B}_2 := L^2(e^{2s\hat{\beta}}\tau^{-5/2}(0, T); L^2(0, L)) \times L^2(0, L)$$

and the operator $\mathcal{A} : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ defined by $\mathcal{A}(z, v) := (z_t + z_{xxx} - \mu(t)z_{xx} + (z\bar{y})_x + zz_x - v1_\omega, z(0))$, for all $(z, v) \in E$.

In order to apply Theorem 5.1, it is necessary to prove that $\mathcal{A}$ is of class $C^1(\mathcal{B}_1, \mathcal{B}_2)$. We start by assuming that $\bar{y} \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L))$. Observe that all terms in the definition of $\mathcal{A}$ are linear (and consequently C^1), except for zz_x . Thus, we will prove that the bilinear operator $((z^1, v^1), (z^2, v^2)) \rightarrow \frac{1}{2}(z^1 z^2)_x$ is continuous from $E \times E$ to $L^2(e^{2s\hat{\beta}}\tau^{-5/2}(0, T); L^2(0, L))$. In fact, notice that

$$e^{s\hat{\beta}}\tau^{-3/2}z \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1(0, L)), \quad \forall (z, v) \in E.$$

Then, we have

$$\begin{aligned} \|e^{2s\hat{\beta}}\tau^{-5/2}(z^1 z^2)_x\|_{L^2(Q)} &\leq C \int_0^T e^{2s\hat{\beta}}\tau^{-3} \|z^1(\cdot, t)\|_{L^\infty(0, L)}^2 e^{2s\hat{\beta}}\tau^{-3} \|z^2(\cdot, t)\|_{H^1(0, L)}^2 \\ &\quad + e^{2s\hat{\beta}}\tau^{-3} \|z^2(\cdot, t)\|_{L^\infty(0, L)}^2 e^{2s\hat{\beta}}\tau^{-3} \|z^1(\cdot, t)\|_{H^1(0, L)}^2 dt \\ &\leq C \|z^1\|_{\mathcal{B}_1} \|z^2\|_{\mathcal{B}_1}. \end{aligned}$$

Now, observe that $\mathcal{A}'(0, 0) : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ is given by

$$\mathcal{A}'(0, 0)(z, v) = (z_t + z_{xxx} - \mu z_{xx} + (z\bar{y})_x - v1_\omega, z(0)), \quad \forall (z, v) \in \mathcal{B}_1.$$

However, the null controllability result proved in Proposition 4.1 allows to deduce that the previous functional is surjective.

Therefore, an application of Theorem 5.1 gives the existence of a positive number δ such that, if $\|z(0)\|_{L^2(0,L)} \leq \delta$, we can find a control v and an associated solution z to (73) satisfying (74). This finishes the proof of Theorem 1.2.

Acknowledgments

This work has been partially supported by FONDECYT grant 1180528 (Eduardo Cerpa), FONDECYT grant 3180100 (Cristhian Montoya), CONICYT–Basal Project FB0008 AC3E (Eduardo Cerpa), Simons Foundation 201615 and NSFC 11571244 (Bingyu Zhang).

References

- [1] V.M. Alekseev, V.M. Tikhomirov, S.V. Fomin, Optimal Control, Contemporary Soviet Mathematics, Consultants Bureau, New York, 1987, translated from the Russian by V.M. Volosov.
- [2] L. Baudouin, E. Cerpa, E. Crépeau, A. Mercado, On the determination of the principal coefficient from boundary measurements in a KdV equation, *J. Inverse Ill-Posed Probl.* 22 (6) (2014) 819–845.
- [3] J.L. Bona, M.E. Schonbek, Travelling-wave solutions to the Korteweg-de Vries-Burgers equation, *Proc. R. Soc. Edinb., Sect. A* 101 (3–4) (1985) 207–226.
- [4] A.-C. Boulanger, P. Trautmann, Sparse optimal control of the KdV-Burgers equation on a bounded domain, *SIAM J. Control Optim.* 55 (6) (2017) 3673–3706.
- [5] J. Boussinesq, Essai sur la théorie des eaux courantes, Mémoires présentés par divers savants à l'Acad. des Sci. Inst. Nat. France 23 (1877) 1–680.
- [6] J.M. Burgers, Application of a model system to illustrate some points of the statistical theory of free turbulence, *Ned. Akad. Wet., Proc.* 43 (1940) 2–12.
- [7] R.A. Capistrano-Filho, A.F. Pazoto, L. Rosier, Internal controllability of the Korteweg–de Vries equation on a bounded domain, *ESAIM Control Optim. Calc. Var.* 21 (4) (2015) 1076–1107.
- [8] M.M. Cavalcanti, V.N. Domingos Cavalcanti, V. Komornik, J.H. Rodrigues, Global well-posedness and exponential decay rates for a KdV-Burgers equation with indefinite damping, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 31 (5) (2014) 1079–1100.
- [9] E. Cerpa, Exact controllability of a nonlinear Korteweg-de Vries equation on a critical spatial domain, *SIAM J. Control Optim.* 46 (3) (2007) 877–899.
- [10] E. Cerpa, E. Crépeau, Boundary controllability for the nonlinear Korteweg-de Vries equation on any critical domain, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 26 (2) (2009) 457–475.
- [11] M. Chen, Bang-bang property for time optimal control of the Korteweg–de Vries–Burgers equation, *Appl. Math. Optim.* 76 (2) (2017) 399–414.
- [12] J.-M. Coron, Control and Nonlinearity, Mathematical Surveys and Monographs, vol. 136, American Mathematical Society, Providence, RI, 2007.
- [13] J.-M. Coron, E. Crépeau, Exact boundary controllability of a nonlinear KdV equation with critical lengths, *J. Eur. Math. Soc.* 6 (3) (2004) 367–398.
- [14] Z. Feng, R. Knobel, Traveling waves to a Burgers-Korteweg-de Vries-type equation with higher-order nonlinearities, *J. Math. Anal. Appl.* 328 (2) (2007) 1435–1450.
- [15] E. Fernández-Cara, S. Guerrero, Null controllability of the Burgers system with distributed controls, *Syst. Control Lett.* 56 (5) (2007) 366–372.
- [16] A. Fursikov, O.Yu. Imanuvilov, Exact controllability of the Navier–Stokes and Boussinesq equations, *Russ. Math. Surv.* 54 (3) (1999) 565–618.
- [17] F.A. Gallego, Controllability aspects of the Korteweg-de Vries Burgers equation on unbounded domains, *J. Math. Anal. Appl.* 461 (1) (2018) 947–970.

- [18] X.-Yi. Gao, Variety of the cosmic plasmas: general variable-coefficient Korteweg-de Vries–Burgers equation with experimental/observational support, *Europhys. Lett.* 110 (1) (2015) 15002.
- [19] O. Glass, S. Guerrero, Some exact controllability results for the linear KdV equation and uniform controllability in the zero-dispersion limit, *Asymptot. Anal.* 60 (1–2) (2008) 61–100.
- [20] S. Guerrero, O.Yu. Imanuvilov, Remarks on global controllability for the Burgers equation with two control forces, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 24 (6) (2007) 897–906.
- [21] S. Guerrero, C. Montoya, Local null controllability of the N -dimensional Navier-Stokes system with nonlinear Navier-slip boundary conditions and $N - 1$ scalar controls, *J. Math. Pures Appl.* (9) 113 (2018) 37–69.
- [22] T. Horsin, On the controllability of the Burgers equation, *ESAIM Control Optim. Calc. Var.* 3 (1998) 83–95.
- [23] C. Jia, Boundary feedback stabilization of the Korteweg–de Vries–Burgers equation posed on a finite interval, *J. Math. Anal. Appl.* 444 (1) (2016) 624–647.
- [24] D.J. Korteweg, G. de Vries, On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves, *Philos. Mag.* (5) 39 (240) (1895) 422–443.
- [25] C. Laurent, L. Rosier, B.-Y. Zhang, Control and stabilization of the Korteweg-de Vries equation on a periodic domain, *Commun. Partial Differ. Equ.* 35 (4) (2010) 707–744.
- [26] J. Li, K. Liu, Well-posedness of Korteweg–de Vries–Burgers equation on a finite domain, *Indian J. Pure Appl. Math.* 48 (1) (2017) 91–116.
- [27] A. Mamun, P. Shukla, The role of dust charge fluctuations on nonlinear dust ion-acoustic waves, *IEEE Trans. Plasma Sci.* 30 (2) (2002) 720–724.
- [28] F. Marbach, Small time global null controllability for a viscous Burgers’ equation despite the presence of a boundary layer, *J. Math. Pures Appl.* (9) 102 (2) (2014) 364–384.
- [29] J.W. Miles, The Korteweg-de Vries equation: a historical essay, *J. Fluid Mech.* 106 (5) (1981) 131–147.
- [30] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York, 1983.
- [31] L. Rosier, Exact boundary controllability for the Korteweg-de Vries equation on a bounded domain, *ESAIM Control Optim. Calc. Var.* 2 (1997) 33–55.
- [32] L. Rosier, Control of the surface of a fluid by a wavemaker, *ESAIM Control Optim. Calc. Var.* 10 (3) (2004) 346–380.
- [33] D.L. Russell, Computational study of the Korteweg-de Vries equation with localized control action, in: G. Chen, E.B. Lee, W. Littman, L. Markus (Eds.), *Distributed Parameter Control Systems: New Trends and Applications*, in: *Lecture Notes in Pure and Appl. Math.*, vol. 128, Marcel Dekker, New York, 1991, pp. 195–203.
- [34] D.L. Russell, B.-Y. Zhang, Exact controllability and stabilizability of the Korteweg-de Vries equation, *Trans. Am. Math. Soc.* 348 (9) (1996) 3643–3672.
- [35] B.-Y. Zhang, *Some Results of Nonlinear Dispersive Wave Equations with Applications to Control*, PhD thesis, University of Wisconsin-Madison, 1990.
- [36] B.-Y. Zhang, Exact boundary controllability of the Korteweg-de Vries equation, *SIAM J. Control Optim.* 37 (2) (1999) 543–565.